

## **PRODUCT INFORMATION**

Tag C-Flag Tag OR2B3 **Target** 

**Synonyms** 6M1-1; OR2B3P; OR6-4; OR6-14

Human OR2B3 full length protein-synthetic Description

nanodisc **Delivery** In Stock **Uniprot ID** 076000 **HEK293 Expression Host** 

**Protein Families Transmembrane** 

**Protein Pathways** Olfactory transduction

The human full length OR2B3 protein has a MW of **Molecular Weight** 

35.5 kDa

Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before Formulation & Reconstitution lyophilization. Please see Certificate of Analysis

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Storage & Shipping

Lyophilized proteins are shipped at ambient

temperature.

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and

hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene

family is the largest in the genome. The nomenclature assigned to the olfactory receptor

genes and proteins for this organism is

independent of other organisms.

Usage Research use only Conjugate Unconjugated

Background

Figure 1. Elisa plates were pre-coated with Flag Tag OR2B3-Nanodisc (0.2μg/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with OR2B3-Nanodisc is 8.804ng/ml. Figure 2. WB analysis of OR2B3-Nanodisc is 8.804ng/ml. Figure 2. WB ana Nanodisc with anti-Flag monoclonal antibody at 1/5000 dilution, followed by Goat Anti-Rabbit IgG HRP at 1/5000 dilution

> Email: info@dimabio.com Website: www.dimabio.com



