

PRODUCT INFORMATION

Tag C-Flag Tag OR8U8 **Target**

Synonyms Olfactory receptor 8U8

Human OR8U8 full length protein-synthetic Description

nanodisc **Delivery** In Stock **Uniprot ID** POC7N1 **Expression Host HEK293**

Protein Families Druggable Genome, Transmembrane

Protein Pathways Olfactory transduction

The human full length OR8U8 protein has a MW of **Molecular Weight**

36.3 kDa

Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before Formulation & Reconstitution lyophilization. Please see Certificate of Analysis

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not Storage & Shipping

intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and

hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor

genes and proteins for this organism is independent of other organisms.

Research use only Usage Conjugate Unconjugated

Background

Figure 1. Elisa plates were pre-coated with Flag Tag OR8U8-Nanodisc (0.2µg/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with OR8U8-Nanodisc is 8.175ng/ml. Figure 2. WB analysis of 0.000 dilutions of the property o Nanodisc with anti-Flag monoclonal antibody at 1/5000 dilution, followed by Goat Anti-Rabbit IgG HRP at 1/5000 dilution

> Email: info@dimabio.com Website: www.dimabio.com

