

PRODUCT INFORMATION

Tag	C-Flag Tag
Target	OXYR
Synonyms	OT-R; OTR
Description	Human OXYR full length protein-synthetic nanodisc
Delivery	In Stock
Uniprot ID	P30559
Expression Host	HEK293
Protein Families	Druggable Genome, GPCR, Transmembrane
Protein Pathways	Calcium signaling pathway, Neuroactive ligand-receptor interaction
Molecular Weight	The human full length OXYR protein has a MW of 42.8 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	The protein belongs to the G-protein coupled receptor family and acts as a receptor for oxytocin. Its activity is mediated by G proteins which activate a phosphatidylinositol-calcium second messenger system. The oxytocin-oxytocin receptor system plays an important role in the uterus during parturition.
Usage	Research use only
Conjugate	Unconjugated



ELISA assay to evaluate OXYR-Nanodisc
0.2 μ g Human OXYR-Nanodisc per well




Figure 1. Elisa plates were pre-coated with Flag Tag OXYR-Nanodisc (0.2 μ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with OXYR-Nanodisc is 11.52ng/ml.




Figure 2. Human OXYR-Nanodisc, Flag Tag on SDS-PAGE

