

PRODUCT INFORMATION

Target	P2RX7
Synonyms	P2X7
Description	Recombinant human P2RX7 protein with N-terminal human Fc tag
Delivery	In Stock
Uniprot ID	Q99572
Expression Host	HEK293
Tag	N-Human Fc Tag
Molecular Characterization	hFc(Glu99-Ala330) P2RX7(Ser47-Val334)
Molecular Weight	The protein has a predicted molecular mass of 59.2 kDa after removal of the signal peptide. The apparent molecular mass of hFc-P2RX7 is approximately 55-70 kDa due to glycosylation.
Purity	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	The product of this gene belongs to the family of purinoreceptors for ATP. This receptor functions as a ligand-gated ion channel and is responsible for ATP-dependent lysis of macrophages through the formation of membrane pores permeable to large molecules. Activation of this nuclear receptor by ATP in the cytoplasm may be a mechanism by which cellular activity can be coupled to changes in gene expression. Multiple alternatively spliced variants have been identified, most of which fit nonsense-mediated decay (NMD) criteria. [provided by RefSeq, Jul 2010]
Usage	Research use only
Conjugate	Unconjugated






Figure 1. Human P2RX7 Protein, hFc Tag on SDS-PAGE under reducing condition.

