

PRODUCT INFORMATION

Target	PCSK9
Synonyms	FH3;FHCL3;HCHOLA3;LDLCQ1;NARC-1;NARC1;PC9
Description	Recombinant Human PCSK9 Protein with C-terminal human Fc tag
Delivery	In Stock
Uniprot ID	Q8NBP7
Expression Host	HEK293
Tag	C-Human Fc Tag
Molecular Characterization	PCSK9(Gln31-Gln692) hFc(Glu99-Ala330)
Molecular Weight	The protein has a predicted molecular mass of 97.2 kDa after removal of the signal peptide. The apparent molecular mass of PCSK9-hFc is approximately 100-130 kDa due to glycosylation.
Purity	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	This gene encodes a member of the subtilisin-like proprotein convertase family, which includes proteases that process protein and peptide precursors trafficking through regulated or constitutive branches of the secretory pathway. The encoded protein undergoes an autocatalytic processing event with its prosegment in the ER and is constitutively secreted as an inactive protease into the extracellular matrix and trans-Golgi network. It is expressed in liver, intestine and kidney tissues and escorts specific receptors for lysosomal degradation. It plays a role in cholesterol and fatty acid metabolism. Mutations in this gene have been associated with autosomal dominant familial hypercholesterolemia. Alternative splicing results in multiple transcript variants.
Usage	Research use only
Conjugate	Unconjugated






Figure 1. Human PCSK9 Protein, hFc Tag on SDS-PAGE under reducing condition.

