

## PRODUCT INFORMATION

<b>Target</b>	PF4V1
<b>Synonyms</b>	CXCL4L1;CXCL4V1;PF4-ALT;PF4A;SCYB4V1
<b>Description</b>	Recombinant Human PF4V1 Protein with C-terminal human Fc tag
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	P10720
<b>Expression Host</b>	HEK293
<b>Tag</b>	C-Human Fc Tag
<b>Molecular Characterization</b>	PF4V1(Phe31-Ser104) hFc(Glu99-Ala330)
<b>Molecular Weight</b>	The protein has a predicted molecular mass of 34.4 kDa after removal of the signal peptide. The apparent molecular mass of PF4V1-hFc is approximately 25-35 kDa due to glycosylation.
<b>Purity</b>	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage&amp;Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	The protein encoded by this gene is a chemokine that is highly similar to platelet factor 4. The encoded protein displays a strong antiangiogenic function and is regulated by chemokine (C-X-C motif) receptor 3. This protein also impairs tumor growth and can protect against blood-retinal barrier breakdown in diabetes patients. [provided by RefSeq, Nov 2015]
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated





Figure 1. Human PF4V1 Protein, hFc Tag on SDS-PAGE under reducing condition.

