

**PRODUCT INFORMATION**

|   |  |
|---|--|
| <b>Tag</b>                              | C-Flag Tag   |
| <b>Target</b>                           | PLLP   |
| <b>Synonyms</b>                         | PMLP, TM4SF11  |
| <b>Description</b>                      | Human PLLP full length protein-synthetic nanodisc  |
| <b>Delivery</b>                         | 6~8weeks   |
| <b>Uniprot ID</b>                       | Q9Y342   |
| <b>Expression Host</b>                  | HEK293   |
| <b>Protein Families</b>                 | Ion Channels: Other  |
| <b>Protein Pathways</b>                 | N/A  |
| <b>Molecular Weight</b>                 | The human full length PLLP protein has a MW of 20kDa   |
| <b>Formulation &amp; Reconstitution</b> | Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.                         |
| <b>Storage&amp;Shipping</b>             | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.                      |
| <b>Background</b>                       | Appears to be involved in myelination. Could also participate in ion transport events as addition of plasmolipin to lipid bilayers induces the formation of ion channels, which are voltage-dependent and K(+)-selective (By similarity).[UniProtKB/Swiss-Prot Function] |
| <b>Usage</b>                            | Research use only  |
| <b>Conjugate</b>                        | Unconjugated   |

