Cat. No. PME100546



## **PRODUCT INFORMATION**

**PSMA Target** 

FGCP;FOLH;GCP2;GCPII;mGCP;NAALAD1;NAALAdase;PSM;PSMA **Synonyms** Recombinant human PSMA protein with N-terminal mouse Fc **Description** 

**Delivery** In Stock **Uniprot ID** 004609 **Expression Host HEK293** 

Tag N-Mouse Fc Tag

Molecular

mFc(Pro99-Lys330) PSMA(Lys44-Ala750) Characterization

The protein has a predicted molecular mass of 105.8 kDa after removal of the signal peptide. The apparent molecular mass of mFc-PSMA is approximately 130 kDa due to glycosylation. **Molecular Weight Purity** 

The purity of the protein is greater than 90% as determined by SDS-PAGE and Coomassie blue staining.

Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please Formulation &

Reconstitution see Certificate of Analysis for specific instructions of

reconstitution.

Storage & Shipping

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. This gene encodes a type II transmembrane glycoprotein belonging to the M28 peptidase family. The protein acts as a glutamate carboxypeptidase on different alternative substrates, including the nutrient folate and the neuropeptide


N-acetyl-l-aspartyl-l-glutamate and is expressed in a number of tissues such as prostate, central and peripheral nervous system and kidney. A mutation in this gene may be associated with impaired intestinal absorption of dietary folates, resulting in low blood folate levels and consequent

**Background** 

In low blood folate levels and consequent hyperhomocysteinemia. Expression of this protein in the brain may be involved in a number of pathological conditions associated with glutamate excitotoxicity. In the prostate the protein is up-regulated in cancerous cells and is used as an effective diagnostic and prognostic indicator of prostate cancer. This gene likely arose from a duplication event of a nearby chromosomal region. Alternative splicing gives rise to multiple transcript variants encoding several different isoforms

Usage Research use only Conjugate Unconjugated

> Email: info@dimabio.com Website: www.dimabio.com










Figure 1. Human PSMA Protein, mFc Tag on SDS-PAGE under reducing condition.

Email: info@dimabio.com Website: www.dimabio.com

