

PRODUCT INFORMATION

C-Flag&Strep Tag Tag

Target PTH1R

Synonyms EKNS; PFE; PTHR; PTHR1

Human PTH1R-Strep full length protein-synthetic **Description**

nanodisc

Delivery In Stock **Uniprot ID** Q03431 **Expression Host HEK293**

Protein Families Druggable Genome, GPCR, Transmembrane

Protein Pathways Neuroactive ligand-receptor interaction

The human full length PTH1R-Strep protein has a **Molecular Weight**

MW of 66.4 kDa Lyophilized from nanodisc solubilization buffer (20

mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Formulation & Reconstitution

for specific instructions. Do not use solvents with a pH below 6.5 or those containing high

concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store Storage & Shipping at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

témperature.

This protein is a receptor for parathyroid hormone (PTH) and for parathyroid hormone-like hormone (PTHLH). The activity of this receptor is mediated by G proteins which activate adenylyl cyclase and

> Email: info@dimabio.com Website: www.dimabio.com

also a phosphatidylinositol-calcium second messenger system. Defects in this receptor are known to be the cause of Jansen's metaphyseal chondrodysplasia (JMC), chondrodysplasia Blomstrand type (BOCD), as well as

enchodromatosis.

Usage Research use only

Conjugate Unconjugated

Background

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China)

/+86-400-006-0995(China)





ELISA assay to evaluate PTH1R-Strep-Nanodisc 0.2µg Human PTH1R-Strep-Nanodisc per well




Figure 1. Elisa plates were pre-coated with C-Flag&Strep Tag PTH1R-Strep-Nanodisc (0.2 μ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with PTH1R-Strep-nanodisc is 1.603ng/ml.



Figure 2. Human PTH1R-Strep-Nanodisc, C-Flag&Strep Tag on SDS-PAGE

Email: info@dimabio.com Website: www.dimabio.com

