

PRODUCT INFORMATION

Tag	C-Flag&Strep Tag
Target	S1PR2
Synonyms	AGR16; DFNB68; EDG-5; EDG5; Gpcr13; H218; LPB2; S1P2
Description	Human S1PR2-Strep full length protein-synthetic nanodisc
Delivery	In Stock
Uniprot ID	O95136
Expression Host	HEK293
Protein Families	Druggable Genome, GPCR, Transmembrane
Protein Pathways	Neuroactive ligand-receptor interaction
Molecular Weight	The human full length S1PR2-Strep protein has a MW of 38.9 kDa Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.
Formulation & Reconstitution	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Storage & Shipping	Receptor for the lysosphingolipid sphingosine 1-phosphate (S1P). S1P is a bioactive lysophospholipid that elicits diverse physiological effect on most types of cells and tissues. When expressed in rat HTC4 hepatoma cells, is capable of mediating S1P-induced cell proliferation and suppression of apoptosis.
Background	
Usage	Research use only
Conjugate	Unconjugated



ELISA assay to evaluate S1PR2-Strep-Nanodisc
0.2µg Human S1PR2-Strep-Nanodisc per well




Figure 1. Elisa plates were pre-coated with C-Flag&Strep Tag S1PR2-Strep-Nanodisc (0.2µg/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with S1PR2-Strep-nanodisc is 1.269ng/ml.




Figure 2. Human S1PR2-Strep-Nanodisc, C-Flag&Strep Tag on SDS-PAGE

