

PRODUCT INFORMATION

| | |
|---|--|
| Target | SCF |
| Synonyms | SF; MGF; SCF; SLF; DCUA; FPH2; FPHH; KL-1; Kitl; WS2F; SHEP7; DFNA69 |
| Description | Recombinant human SCF Protein with C-terminal 10×His tag |
| Delivery | In Stock |
| Uniprot ID | P21583 |
| Expression Host | HEK293 |
| Tag | C-10×His tag |
| Molecular Characterization | SCF(Glu26-His214) 10×His tag |
| Molecular Weight | The protein has a predicted molecular mass of 22.4.0 kDa after removal of the signal peptide. The purity of the protein is greater than 85% as determined by SDS-PAGE and Coomassie blue staining. |
| Purity | Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. |
| Formulation & Reconstitution | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Storage&Shipping | This gene encodes the ligand of the tyrosine-kinase receptor encoded by the KIT locus. This ligand is a pleiotropic factor that acts in utero in germ cell and neural cell development, and hematopoiesis, all believed to reflect a role in cell migration. In adults, it functions pleiotropically, while mostly noted for its continued requirement in hematopoiesis. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008] |
| Background | Research use only |
| Usage | Unconjugated |






Figure 1. Human SCF Protein, His Tag on SDS-PAGE under reducing condition.

