

PRODUCT INFORMATION

Tag	C-Flag&Strep Tag
Target	SCTR
Synonyms	SR
Description	Human SCTR-Strep full length protein-synthetic nanodisc
Delivery	In Stock
Uniprot ID	P47872
Expression Host	HEK293
Protein Families	Druggable Genome, GPCR, Transmembrane
Protein Pathways	Neuroactive ligand-receptor interaction
Molecular Weight	The human full length SCTR-Strep protein has a MW of 50.2 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	G protein-coupled receptor activated by secretin (SCT), which is involved in different processes such as regulation of the pH of the duodenal content, food intake and water homeostasis. It binds secretin which is the most potent regulator of pancreatic bicarbonate, electrolyte and volume secretion. Secretin and its receptor are suggested to be involved in pancreatic cancer and autism.
Usage	Research use only
Conjugate	Unconjugated



**ELISA assay to evaluate SCTR-Strep-Nanodisc**  
**0.2µg Human SCTR-Strep-Nanodisc per well**




Figure 1. Elisa plates were pre-coated with C-Flag&Strep Tag SCTR-Strep-Nanodisc (0.2µg/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with SCTR-Strep-nanodisc is 1.208ng/ml.




Figure 2. Human SCTR-Strep-Nanodisc, C-Flag&Strep Tag on SDS-PAGE

