

PRODUCT INFORMATION

Tag	C-Flag Tag
Target	SLC25A4
Synonyms	AAC1; ANT; ANT 1; ANT1; MTDPS12; MTDPS12A; PEO2; PEO3; PEOA2; T1
Description	Human SLC25A4 full length protein-synthetic nanodisc
Delivery	In Stock
Uniprot ID	P12235
Expression Host	HEK293
Protein Families	Druggable Genome, Transmembrane
Protein Pathways	Calcium signaling pathway, Huntington's disease, Parkinson's disease
Molecular Weight	The human full length SLC25A4 protein has a MW of 33.1 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	The protein functions as a gated pore that translocates ADP from the cytoplasm into the mitochondrial matrix and ATP from the mitochondrial matrix into the cytoplasm. The protein forms a homodimer embedded in the inner mitochondria membrane. Mutations in this gene have been shown to result in autosomal dominant progressive external ophthalmoplegia and familial hypertrophic cardiomyopathy.
Usage	Research use only
Conjugate	Unconjugated






Figure1. Elisa plates were pre-coated with Flag Tag SLC25A4-Nanodisc (0.2µg/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with SLC25A4-Nanodisc is 36.85ng/ml.




Figure2. Human SLC25A4-Nanodisc, Flag Tag on SDS-PAGE

