

## PRODUCT INFORMATION

<b>Tag</b>	C-Flag Tag
<b>Target</b>	SLC2A4
<b>Synonyms</b>	GLUT4
<b>Description</b>	Human SLC2A4 full length protein-synthetic nanodisc
<b>Delivery</b>	3-4 weeks
<b>Uniprot ID</b>	P14672
<b>Expression Host</b>	HEK293
<b>Protein Families</b>	Druggable Genome, Transmembrane
<b>Protein Pathways</b>	Adipocytokine signaling pathway, Insulin signaling pathway, Type II diabetes mellitus
<b>Molecular Weight</b>	The human full length SLC2A4 protein has a MW of 54.6 kDa
<b>Formulation &amp; Reconstitution</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Storage&amp;Shipping</b>	This gene is a member of the solute carrier family 2 (facilitated glucose transporter) family and encodes a protein that functions as an insulin-regulated facilitative glucose transporter. In the absence of insulin, this integral membrane protein is sequestered within the cells of muscle and adipose tissue. Within minutes of insulin stimulation, the protein moves to the cell surface and begins to transport glucose across the cell membrane. Mutations in this gene have been associated with noninsulin-dependent diabetes mellitus (NIDDM). [provided by RefSeq, Jul 2008]
<b>Background</b>	Research use only
<b>Usage</b>	Unconjugated
<b>Conjugate</b>	



**ELISA assay to evaluate SLC2A4-Nanodisc**  
0.2 $\mu$ g Human SLC2A4-Nanodisc per well




Figure1. Elisa plates were pre-coated with Flag Tag SLC2A4-Nanodisc (0.2 $\mu$ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with SLC2A4-Nanodisc is 4.357ng/ml.



Figure2. Human SLC2A4-Nanodisc, Flag Tag on SDS-PAGE

