

**PRODUCT INFORMATION**

|   |  |
|---|--|
| <b>Tag</b>                              | C-Flag&Strep Tag   |
| <b>Target</b>                           | SLC4A3   |
| <b>Synonyms</b>                         | AE3; CAE3/BAE3; SLC2C  |
| <b>Description</b>                      | Human SLC4A3-Strep full length protein-synthetic nanodisc  |
| <b>Delivery</b>                         | In Stock   |
| <b>Uniprot ID</b>                       | P48751   |
| <b>Expression Host</b>                  | HEK293   |
| <b>Protein Families</b>                 | Transmembrane  |
| <b>Protein Pathways</b>                 | N/A  |
| <b>Molecular Weight</b>                 | The human full length SLC4A3-Strep protein has a MW of 135.8 kDa   |
| <b>Formulation &amp; Reconstitution</b> | Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments. |
| <b>Storage&amp;Shipping</b>             | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.  |
| <b>Background</b>                       | The protein is a plasma membrane anion exchange protein. The encoded protein has been found in brain, heart, kidney, small intestine, and lung.  |
| <b>Usage</b>                            | Research use only  |
| <b>Conjugate</b>                        | Unconjugated   |



### ELISA assay to evaluate SLC4A3-Strep-Nanodisc 0.2 $\mu$ g Human SLC4A3-Strep-Nanodisc per well




Figure 1. Elisa plates were pre-coated with C-Flag&Strep Tag SLC4A3-Strep-Nanodisc (0.2 $\mu$ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with SLC4A3-Strep-nanodisc is 4.136ng/ml.



Figure 2. Human SLC4A3-Strep-Nanodisc, C-Flag&Strep Tag on SDS-PAGE

