

## **PRODUCT INFORMATION**

ST2 **Target** 

T1; IL1RL1; DER4; ST2L; ST2V; FIT-1; IL33R **Synonyms** 

Recombinant human ST2 Protein with C-terminal **Description** 

6×His tag

**Delivery** In Stock **Uniprot ID** Q01638 **Expression Host HEK293** 

Tag C-6×His tag

Molecular Characterization

Storage & Shipping

**Background** 

Purity

ST2(Lys19-Ser328) 6×His tag

The protein has a predicted molecular mass of **Molecular Weight** 

35.8 kDa after removal of the signal peptide. The apparent molecular mass of ST2-His is

approximately 35-70 kDa due to glycosylation. The purity of the protein is greater than 85% as determined by SDS-PAGE and Coomassie blue

staining.

Lyophilized from sterile PBS, pH 7.4. Normally 5 % 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Formulation & Reconstitution

for specific instructions.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store

at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

The protein encoded by this gene is a member of the interleukin 1 receptor family. Studies of the similar gene in mouse suggested that this receptor can be induced by proinflammatory stimuli, and may be involved in the function of helper T cells. This gene, interleukin 1 receptor, type I (IL1R1), interleukin 1 receptor, type II (IL1R1)

(IL1R2) and interleukin 1 receptor-like 2 (IL1RL2) form a cytokine receptor gene cluster in a region mapped to chromosome 2q12. Alternative splicing of this gene results in multiple transcript variants. [provided by RefSeq, Jul 2008]

Research use only Usage Conjugate Unconjugated










Figure 1. Human ST2 Protein, His Tag on SDS-PAGE under reducing condition.

## Human ST2, His Tagged protein ELISA

0.2 μg of Human ST2, His tagged protein per well




Figure 2. ELISA plate pre-coated by 2  $\mu$ g/mL (100  $\mu$ L/well) Human ST2 Protein, His Tag (PME101359) can bind Anti-ST2(astegolimab biosimilar) mAb (BME100263) in a linear range of 0.64–80 ng/mL.

Email: info@dimabio.com Website: www.dimabio.com

