

PRODUCT INFORMATION

| | |
|---|--|
| Tag | C-Flag&Strep Tag |
| Target | TA2R9 |
| Synonyms | T2R9, TRB6 |
| Description | Human TA2R9-Strep full length protein-synthetic nanodisc |
| Delivery | 6~8weeks |
| Uniprot ID | Q9NYW1 |
| Expression Host | HEK293 |
| Protein Families | Transmembrane,Druggable Genome, |
| Protein Pathways | N/A |
| Molecular Weight | The human full length TA2R9-Strep protein has a MW of 35.6 kDa |
| Formulation & Reconstitution | Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments. |
| Storage&Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | This gene product belongs to the family of candidate taste receptors that are members of the G-protein-coupled receptor superfamily. These proteins are specifically expressed in the taste receptor cells of the tongue and palate epithelia. They are organized in the genome in clusters and are genetically linked to loci that influence bitter perception in mice and humans. In functional expression studies, they respond to bitter tastants. This gene maps to the taste receptor gene cluster on chromosome 12p13. [provided by RefSeq, Jul 2008] |
| Usage | Research use only |
| Conjugate | Unconjugated |

