

## **PRODUCT INFORMATION**

**Target** TGFB1

CED; LAP; DPD1; TGFB; IBDIMDE; TGFbeta; TGF-**Synonyms** 

beta1

Recombinant human TGFB1(30-390) Protein with **Description** 

C-terminal 10×His tag

Delivery In Stock P01137 **Uniprot ID Expression Host** HFK293

Tag C-10×His tag

Molecular

Storage & Shipping

Background

**Purity** 

TGFB1(Leu30-Ser390) 10×His tag Characterization

The protein has a predicted molecular mass of **Molecular Weight** 42.6 kDa after removal of the signal peptide.

The purity of the protein is greater than 80% as determined by SDS-PAGE and Coomassie blue

Lyophilized from sterile PBS, pH 7.4. Normally 5 % Formulation & - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Reconstitution

for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store

at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

This gene encodes a secreted ligand of the TGF-beta (transforming growth factor-beta) superfamily of proteins. Ligands of this family

bind various TGF-beta receptors leading to recruitment and activation of SMAD family transcription factors that regulate gene expression. The encoded preproprotein is proteolytically processed to generate a latency-associated peptide (LAP) and a mature peptide, and is found in either a latent form composed of a mature peptide homodimer, a LAP homodimer,

and a latent TGF-beta binding protein, or in an active form consisting solely of the mature peptide homodimer. The mature peptide may also form heterodimers with other TGFB family members. This encoded protein regulates cell proliferation, differentiation and growth, and can modulate expression and activation of other growth factors including interferon gamma and tumor necrosis factor alpha. This gene is frequently upregulated in tumor cells, and mutations in this gene result in Camurati-

> Email: info@dimabio.com Website: www.dimabio.com

Engelmann disease. [provided by RefSeq, Aug 2016]

**Usage** Research use only

Conjugate Unconjugated

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)






Figure 1. Human TGFB1(30-390) Protein, His Tag on SDS-PAGE under reducing condition.

## Human TGFB1(30-390), His Tagged protein ELISA

0.2 μg of Human TGFB1(30-390), His tagged protein per well




Figure 2. ELISA plate pre-coated by 2  $\mu$ g/mL (100  $\mu$ L/well) Human TGFB1(30-390) Protein, His Tag (PME101649) can bind Human TGFBR2 Protein, hFc Tag (PME101017) in a linear range of 3.20–400 ng/mL.

Email: info@dimabio.com Website: www.dimabio.com

