

## PRODUCT INFORMATION

<b>Target</b>	TGFBR1
<b>Synonyms</b>	AAT5;ACVRLK4;ALK-5;ALK5;ESS1;LDS1;LDS1A;LDS2A;MSSE;SKR4;tbetaR-I;TBR-i;TBRI;TGFR-1
<b>Description</b>	Recombinant human TGFBR1 Protein with N-terminal Human Fc tag
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	P36897
<b>Expression Host</b>	HEK293
<b>Tag</b>	N-Human Fc Tag
<b>Molecular Characterization</b>	hFc(Glu99-Ala330) TGFBR1(Leu34-Leu126)
<b>Molecular Weight</b>	The protein has a predicted molecular mass of 36.3 kDa after removal of the signal peptide. The apparent molecular mass of hFc-TGFBR1 is approximately 40-53 kDa due to glycosylation.
<b>Purity</b>	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage&amp;Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	The protein encoded by this gene forms a heteromeric complex with type II TGF-beta receptors when bound to TGF-beta, transducing the TGF-beta signal from the cell surface to the cytoplasm. The encoded protein is a serine/threonine protein kinase. Mutations in this gene have been associated with Loeys-Dietz aortic aneurysm syndrome (LDAS). Multiple transcript variants encoding different isoforms have been found for this gene.
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated



Figure 1. Human TGFBR1 Protein, N-hFc Tag on SDS-PAGE under reducing condition.

