**Delivery** 

**Background** 



## **PRODUCT INFORMATION**

C-Flag&Strep Tag Tag

TLR4 **Target** 

**Synonyms** ARMD10; CD284; TLR-4; TOLL

Human TLR4-Strep full length protein-synthetic **Description** 

nanodisc In Stock

**Uniprot ID** 000206 **Expression Host HEK293** 

Druggable Genome, Transmembrane **Protein Families Protein Pathways** Toll-like receptor signaling pathway

The human full length TLR4-Strep protein has a **Molecular Weight** 

MW of 95.7 kDa

Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before Formulation & Reconstitution lyophilization. Please see Certificate of Analysis

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Storage & Shipping

Lyophilized proteins are shipped at ambient

temperature.

The protein is a member of the Toll-like receptor (TLR) family which plays a fundamental role in pathogen recognition and activation of innate immunity. TLRs are highly conserved from Drosophila to humans and share structural and functional similarities. They recognize pathogenassociated molecular patterns that are expressed on infectious agents, and mediate the production of cytokines necessary for the development of effective immunity. The various TLRs exhibit different patterns of expression. In silico studies have found a particularly strong binding of

surface TLR4 with the spike protein of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), the causative agent of Coronavirus disease-2019 (COVID-19). This receptor has also been implicated in signal transduction events induced by lipopolysaccharide (LPS) found in most gram-negative bacteria. Mutations in this gene have been associated with differences in LPS responsiveness, and with susceptibility to

age-related macular degeneration.

**Usage** Research use only Conjugate Unconjugated

Figure 1. Elisa plates were pre-coated with C-Flag&Strep Tag TLR4-Strep-Nanodisc (0.2µg/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with TLR4-Strep-nanodisc is 2.447ng/ml. Email: info@dimabio.com

Website: www.dimabio.com

Aress: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-1806274953(China) /+86-400-006-0995(China)

Human TLR4-Strep full length protein-synthetic nanodisc

Cat. No. FLP120141



Figure 2. Human TLR4-Strep-Nanodisc, C-Flag&Strep Tag on SDS-PAGE

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Email: info@dimabio.com Website: www.dimabio.com

