

PRODUCT INFORMATION

TNFRSF10B **Target**

Synonyms TNFRSF10B;TRAILR2;TRAIL-R2;CD262;DR5;KILLER;TRICK2;ZTNFR9;TRICKB Description Recombinant human TNFRSF10B protein with C-terminal mouse Fc tag

Delivery In Stock **Uniprot ID** 014763 **HEK293 Expression Host**

C-Mouse Fc Tag

Molecular Characterization

TNFRSF10B (Ile56-Glu182) mFc(Pro99-Lys330)

The protein has a predicted molecular mass of 40.5 kDa after removal of the signal peptide. $\label{eq:protein} % \begin{center} \begin{cen$ **Molecular Weight**

The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining. **Purity**

Formulation & Reconstitution

Storage & Shipping

and Coomassie blue staining.

Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.

The protein encoded by this gene is a member of the TNF-receptor superfamily, and contains an intracellular death domain. This receptor can be activated by tumor necrosis factor-related apoptosis inducing ligand (TNFSF10/TRAIL/APO-2L), and transduces an apoptosis signal. Studies with FADD-deficient mice suggested that FADD, a death domain containing adaptor protein, is required for the apoptosis mediated by this protein. Two transcript variants encoding different isoforms and one noncoding transcript have been found for this gene.

Research use only

Usage Research use only Unconjugated Conjugate

Background



Email: info@dimabio.com








Figure 1. Human TNFRSF10B Protein, mFc Tag on SDS-PAGE under reducing condition.

Human TNFRSF10B, mFc Tagged protein ELISA

0.2 µg of Human TNFRSF10B, mFc tagged protein per well




Figure 2.ELISA plate pre-coated by 2 μ g/ml (100 μ l/well) Human TNFRSF10B, mFc tagged protein (PME100465) can bind Anti-TNFRSF10B Neutralizing antibody ([getskuurl sku="BME100032"]) in a linear range of 3.2-80 ng/ml.

Email: info@dimabio.com Website: www.dimabio.com

