

PRODUCT INFORMATION

TPO **Target**

Thrombopoietin; C-mpl ligand; Megakaryocyte colony-stimulating factor;Megakaryocyte growth and development factor;Myeloproliferative leukemia virus oncogene ligand;THPO **Synonyms**

Recombinant Human Thrombopoietin is produced by our Mammalian expression system and the target gene encoding Ser22-Gly353 is expressed with a 6His tag at the N-terminus, 6His tag at the

C-terminus.

Delivery In Stock P40225 **Uniprot ID Expression Host HEK293**

N, C-6×His Tag Tag

Molecular

Storage & Shipping

Background

Description

Not available Characterization

37.3 KDa Molecular Weight

Greater than 95% as determined by reducing **Purity**

SDS-PAGE.

Lyophilized from a 0.2 µm filtered solution of 20mM Tris-HCl, 150mM NaCl, pH 8.0. Formulation &

Reconstitution

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

temperature.

Thrombopoietin (TPO) is a glycoprotein hormone which belongs to the EPO/TPO family. It produced

by the liver and kidney which regulates the production of platelets. TPO stimulates the production and differentiation of megakaryocytes, the bone marrow cells that bud off large numbers of platelets. Lineage-specific cytokine affects the proliferation and maturation of megakaryocytes from their committed progenitor cells. It acts at a late stage of megakaryocyte development. It may

late stage of megakaryocyte development. It may be the major physiological regulator of circulating

> Email: info@dimabio.com Website: www.dimabio.com

platelets.

Usage Research use only

Conjugate Unconjugated








Figure 1. Greater than 95% as determined by reducing SDS-PAGE.

Email: info@dimabio.com Website: www.dimabio.com

