

## PRODUCT INFORMATION

<b>Tag</b>	C-Flag&Strep Tag
<b>Target</b>	TRPM5
<b>Synonyms</b>	LTRPC5, MTR1
<b>Description</b>	Human TRPM5-Strep full length protein-synthetic nanodisc
<b>Delivery</b>	6~8weeks
<b>Uniprot ID</b>	Q9NZQ8
<b>Expression Host</b>	HEK293
<b>Protein Families</b>	Ion Channels: Transient receptor potential
<b>Protein Pathways</b>	N/A
<b>Molecular Weight</b>	The human full length TRPM5-Strep protein has a MW of 131.5 kDa
<b>Formulation &amp; Reconstitution</b>	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage&amp;Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	This gene encodes a member of the transient receptor potential (TRP) protein family, which is a diverse group of proteins with structural features typical of ion channels. This protein plays an important role in taste transduction, and has characteristics of a calcium-activated, non-selective cation channel that carries Na <sup>+</sup> , K <sup>+</sup> , and Cs <sup>+</sup> ions equally well, but not Ca <sup>2+</sup> ions. It is activated by lower concentrations of intracellular Ca <sup>2+</sup> , and inhibited by higher concentrations. It is also a highly temperature-sensitive, heat activated channel showing a steep increase of inward currents at temperatures between 15 and 35 degrees Celsius. This gene is located within the Beckwith-Wiedemann syndrome critical region-1 on chromosome 11p15.5, and has been shown to be imprinted, with exclusive expression from the paternal allele. [provided by RefSeq, Oct 2010]
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated

