

PRODUCT INFORMATION

Tag C-Flag&Strep Tag

Target AVPR2

Synonyms ADHR; DI1; DIR; DIR3; NDI; NDI1; V2R

DescriptionHuman AVPR2-Strep full length protein-synthetic

nanodisc

Delivery In Stock
Uniprot ID P30518
Expression Host HEK293

Formulation & Reconstitution

Storage & Shipping

Background

Protein Families Druggable Genome, GPCR, Transmembrane

Protein Pathways Neuroactive ligand-receptor interaction

Molecular Weight

The human full length AVPR2-Strep protein has a

MW of 40.3 kDa

mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with

Lyophilized from nanodisc solubilization buffer (20

for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

temperature.

Vasopressin receptor type 2, also known as the V2 receptor, which belongs to the seven-transmembrane-domain G protein-coupled receptor (GPCR) superfamily, and couples to Gs thus stimulating adenylate cyclase. The subfamily that includes the V2 receptor, the V1a and V1b vasopressin receptors, the oxytocin receptor, and

isotocin and mesotocin receptors in nonmammals, is well conserved, though several members signal via other G proteins. All bind similar cyclic nonapeptide hormones. The V2 receptor is expressed in the kidney tubule, predominantly in the distal convoluted tubule and collecting ducts, where its primary property is to

respond to the pituitary hormone arginine vasopressin (AVP) by stimulating mechanisms that concentrate the urine and maintain water homeostasis in the organism. When the function of this gene is lost, the disease Nephrogenic Diabetes Insipidus (NDI) results. The V2 receptor is also expressed outside the kidney although its tissue localization is uncertain. When these 'extrarenal receptors' are stimulated by infusion of a V2 selective agonist (dDAVP), a variety of clotting factors are released into the bloodstream. The physiologic importance of this property is not known - its absence does not appear to be

known – its absence does not appear to be detrimental in NDI patients. The gene expression has also been described in fetal lung tissue and lung cancer associated with alternative splicing

Email: info@dimabio.com

Website: www.dimabio.com

lung cancer associated with alternative splicing.

Usage Research use only

Conjugate Unconjugated

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)



ELISA assay to evaluate AVPR2-Strep-Nanodisc 0.2µg Human AVPR2-Strep-Nanodisc per well




Figure 1. Elisa plates were pre-coated with C-Flag&Strep Tag AVPR2-Strep-Nanodisc (0.2 μ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with AVPR2-Strep-nanodisc is 2.798ng/ml.



Figure 2. Human AVPR2-Strep-Nanodisc, C-Flag&Strep Tag on SDS-PAGE
Address: Wuhan institute of Email: info@dimabio.com
Biotechnology B7, Biolake No.666 Website: www.dimabio.com

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China) Human AVPR2-Strep full length protein-synthetic nanodisc

Cat. No. FLP120492





