

PRODUCT INFORMATION

C-Flag&Strep Tag Tag

VDAC1 **Target**

Synonyms PORIN; VDAC-1

Human VDAC1-Strep full length protein-synthetic **Description**

nanodisc **Delivery** In Stock **Uniprot ID** P21796

Expression Host HEK293

Formulation & Reconstitution

Background

Protein Families Ion Channels: Other

Calcium signaling pathway, Huntington's disease, **Protein Pathways**

Parkinson's disease

The human full length VDAC1-Strep protein has a **Molecular Weight**

MW of 30.8 kDa

Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis

for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store

Storage & Shipping at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient


temperature.

A voltage-dependent anion channel protein that is a major component of the outer mitochondrial membrane. The encoded protein facilitates the exchange of metabolites and ions across the outer mitochondrial membrane and may regulate mitochondrial functions. This protein also forms channels in the plasma membrane and may be

involved in transmembrane electron transport. Alternate splicing results in multiple transcript variants. Multiple pseudogenes of this gene are found on chromosomes 1, 2 3, 6, 9, 12, X and Y.

Usage Research use only Conjugate Unconjugated

Email: info@dimabio.com Website: www.dimabio.com







ELISA assay to evaluate VDAC1-Strep-Nanodisc 0.2µg Human VDAC1-Strep-Nanodisc per well




Figure 1. Elisa plates were pre-coated with C-Flag&Strep Tag VDAC1-Strep-Nanodisc (0.2 μ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with VDAC1-Strep-nanodisc is 5.450ng/ml.



Figure 2. Human VDAC1-Strep-Nanodisc, C-Flag&Strep Tag on SDS-PAGE

Email: info@dimabio.com Website: www.dimabio.com

