

PRODUCT INFORMATION

Tag	C-Flag&Strep Tag
Target	VDAC3
Synonyms	HD-VDAC3, VDAC-3
Description	Human VDAC3-Strep full length protein-synthetic nanodisc
Delivery	6~8weeks
Uniprot ID	Q9Y277
Expression Host	HEK293
Protein Families	Ion Channels: Other
Protein Pathways	N/A
Molecular Weight	The human full length VDAC3-Strep protein has a MW of 30.7 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	This gene encodes a voltage-dependent anion channel (VDAC), and belongs to the mitochondrial porin family. VDACS are small, integral membrane proteins that traverse the outer mitochondrial membrane and conduct ATP and other small metabolites. They are known to bind several kinases of intermediary metabolism, thought to be involved in translocation of adenine nucleotides, and are hypothesized to form part of the mitochondrial permeability transition pore, which results in the release of cytochrome c at the onset of apoptotic cell death. Alternatively transcript variants encoding different isoforms have been described for this gene. [provided by RefSeq, Oct 2011]
Usage	Research use only
Conjugate	Unconjugated

