

## PRODUCT INFORMATION

|                                   |  |
|-----------------------------------|--|
| <b>Target</b>                     | 4-1BB Ligand   |
| <b>Synonyms</b>                   | Tnfsf9   |
| <b>Description</b>                | Recombinant mouse 4-1BB Ligand protein with N-terminal human Fc tag  |
| <b>Delivery</b>                   | In Stock   |
| <b>Uniprot ID</b>                 | P41274   |
| <b>Expression Host</b>            | HEK293   |
| <b>Tag</b>                        | N-Human Fc Tag   |
| <b>Molecular Characterization</b> | <p>hFc(Glu99-Ala330) Mouse 4-1BB Ligand(Arg104-Glu309)</p> <p>The protein has a predicted molecular mass of 49.1 kDa after removal of the signal peptide. The apparent molecular mass of hFc-m4-1BB Ligand is approximately 55-70 kDa due to glycosylation.</p>  |
| <b>Molecular Weight</b>           | <p>The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.</p> <p>Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.</p>  |
| <b>Storage&amp;Shipping</b>       | <p>Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).</p> <p>Lyophilized proteins are shipped at ambient temperature.</p> <p>Cytokine that binds to TNFRSF9. Induces the proliferation of activated peripheral blood T-cells. May have a role in activation-induced cell death (AICD). May play a role in cognate interactions between T-cells and B-cells/macrophages.[UniProtKB/Swiss-Prot Function]</p> |
| <b>Background</b>                 | Research use only  |
| <b>Usage</b>                      | Research use only  |
| <b>Conjugate</b>                  | Unconjugated   |






Figure 1. Mouse 4-1BB Ligand Protein, hFc Tag on SDS-PAGE under reducing condition.

