

PRODUCT INFORMATION

| | |
|---|---|
| Target | CD27 |
| Synonyms | S152;Tnfrsf7;Tp55 |
| Description | Recombinant mouse CD27 protein with C-terminal human Fc tag |
| Delivery | In Stock |
| Uniprot ID | P41272 |
| Expression Host | HEK293 |
| Tag | C-Human Fc Tag |
| Molecular Characterization | Mouse CD27(Pro24-Arg182) hFc(Glu99-Ala330) |
| Molecular Weight | The protein has a predicted molecular mass of 43.9 kDa after removal of the signal peptide. The apparent molecular mass of mCD27-hFc is approximately 55-70 kDa due to glycosylation. |
| Purity | The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining. |
| Formulation & Reconstitution | Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. |
| Storage&Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | Receptor for CD70/CD27L. May play a role in survival of activated T-cells. May play a role in apoptosis through association with SIVA1.[UniProtKB/Swiss-Prot Function] |
| Usage | Research use only |
| Conjugate | Unconjugated |






Figure 1. Mouse CD27 Protein, hFc Tag on SDS-PAGE under reducing condition.

