

## PRODUCT INFORMATION

<b>Tag</b>	C-Flag&Strep Tag
<b>Target</b>	CLDN18.1
<b>Synonyms</b>	N/A
<b>Description</b>	Mouse Cldn18.1-Strep full length protein-synthetic nanodisc
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	P56857-1
<b>Expression Host</b>	HEK293
<b>Protein Families</b>	N/A
<b>Protein Pathways</b>	N/A
<b>Molecular Weight</b>	The mouse full length Cldn18.1-Strep protein has a MW of 28.1 kDa Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Formulation &amp; Reconstitution</b>	A member of the claudin family. Claudins are integral membrane proteins and components of tight junction strands. Tight junction strands serve as a physical barrier to prevent solutes and water from passing freely through the paracellular space between epithelial or endothelial cell sheets, and also play critical roles in maintaining cell polarity and signal transductions. This gene is a downstream target gene regulated by the T/EBP/NKX2.1 homeodomain transcription factor. Four alternatively spliced transcript variants resulted from alternative promoters and alternative splicing have been identified, which encode two lung-specific isoforms and two stomach-specific isoforms respectively. This gene is also expressed in colons, inner ear and skin, and its expression is increased in both experimental colitis and ulcerative colitis.
<b>Storage&amp;Shipping</b>	Research use only
<b>Background</b>	Unconjugated
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated




**ELISA assay to evaluate Cldn18.1-Strep-Nanodisc**
0.2 $\mu$ g mouse Cldn18.1-Strep-Nanodisc per well

Figure 1. Elisa plates were pre-coated with C-Flag&Strep Tag Cldn18.1-Strep-Nanodisc (0.2 $\mu$ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with Cldn18.1-Strep-nanodisc is 4.205ng/ml.




Figure 2. Mouse Cldn18.1-Strep-Nanodisc, C-Flag&Strep Tag on SDS-PAGE

