

PRODUCT INFORMATION

| | |
|---|--|
| Clone ID | DMC682 |
| Target | CD166 |
| Synonyms | CD166; MEMD |
| Host Species | Rabbit |
| Description | PE-conjugated Anti-CD166 antibody(DMC682), IgG1 Chimeric mAb |
| Delivery | Under Development |
| Uniprot ID | Q13740-1 |
| IgG type | Rabbit/Human Fc chimeric IgG1 |
| Clonality | Monoclonal |
| Reactivity | Human |
| Applications | Flow Cyt |
| Recommended Dilutions | Flow Cyt 1:100 |
| Purification | Purified from cell culture supernatant by affinity chromatography |
| Formulation & Reconstitution | Liquid PBS with 0.05% Proclin300, 1% BSA |
| Storage&Shipping | Store at 2°C-8°C for 6 months |
| Background | This gene encodes activated leukocyte cell adhesion molecule (ALCAM); also known as CD166 (cluster of differentiation 166); which is a member of a subfamily of immunoglobulin receptors with five immunoglobulin-like domains (VVC2C2C2) in the extracellular domain. This protein binds to T-cell differentiation antigen CD6; and is implicated in the processes of cell adhesion and migration. Multiple alternatively spliced transcript variants encoding different isoforms have been found. [provided by RefSeq; Aug 2011] |
| Usage | Research use only |
| Conjugate | PE-conjugated |
| DIMA Disclaimer | All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scr |

