

**PRODUCT INFORMATION**

<b>Clone ID</b>	DM155
<b>Target</b>	CD171
<b>Synonyms</b>	CAML1; CD171; HSAS; HSAS1; MASA; MIC5; N-CAM-L1; N-CAML1; NCAM-L1; S10; SPG1
<b>Host Species</b>	Rabbit
<b>Description</b>	PE-conjugated Anti-CD171 antibody(DM155); Rabbit mAb
<b>Delivery</b>	Under Development
<b>Uniprot ID</b>	P32004
<b>IgG type</b>	Rabbit IgG
<b>Clonality</b>	Monoclonal
<b>Reactivity</b>	Human
<b>Applications</b>	Flow Cyt
<b>Recommended Dilutions</b>	Flow Cyt 1:100
<b>Purification</b>	Purified from cell culture supernatant by affinity chromatography
<b>Endotoxin</b>	Less than 1.0 EU/μg by the LAL method. For <1 EU/mg requirements, please contact us for customization.
<b>Formulation &amp; Reconstitution</b>	Liquid□PBS with 0.05% Proclin300, 1% BSA
<b>Storage&amp;Shipping</b>	Store at 2°C-8°C for 6 months
<b>Background</b>	The protein encoded by this gene is an axonal glycoprotein belonging to the immunoglobulin supergene family. The ectodomain; consisting of several immunoglobulin-like domains and fibronectin-like repeats (type III); is linked via a single transmembrane sequence to a conserved cytoplasmic domain. This cell adhesion molecule plays an important role in nervous system development; including neuronal migration and differentiation. Mutations in the gene cause X-linked neurological syndromes known as CRASH (corpus callosum hypoplasia; retardation; aphasia; spastic paraplegia and hydrocephalus). Alternative splicing of this gene results in multiple transcript variants; some of which include an alternate exon that is considered to be specific to neurons.
<b>Usage</b>	Research use only
<b>Conjugate</b>	PE-conjugated
<b>DIMA Disclaimer</b>	All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scr

