

PRODUCT INFORMATION

| | |
|---|--|
| Clone ID | 19A5 |
| Target | FLT3 |
| Synonyms | FIt-3;Flk-2;STK-1;CD135;FLK2;FLT-3 |
| Host Species | Rabbit |
| Description | PE-conjugated Anti-FLT3 antibody(19A5), IgG1 Chimeric mAb |
| Delivery | Under Development |
| Uniprot ID | P36888 |
| IgG type | Rabbit/Human Fc chimeric IgG1 |
| Clonality | Monoclonal |
| Reactivity | Human |
| Applications | Flow Cyt |
| Recommended Dilutions | Flow Cyt 1:100 |
| Purification | Purified from cell culture supernatant by affinity chromatography |
| Formulation & Reconstitution | Liquid□PBS with 0.05% Proclin300, 1% BSA |
| Storage&Shipping | Store at 2°C-8°C for 6 months |
| Background | <p>FIt-3(Receptor-type tyrosine-protein kinase FLT3) is also known as FLK-2(Fetal liver kinase-2), STK-1(Stem cell tyrosine kinase 1), CD135. FLT3 is a cytokine receptor which belongs to the receptor tyrosine kinase class III. Tyrosine-protein kinase that acts as cell-surface receptor for the cytokine FLT3LG and regulates differentiation, proliferation and survival of hematopoietic progenitor cells and of dendritic cells. Promotes phosphorylation of SHC1 and AKT1, and activation of the downstream effector MTOR. Promotes activation of RAS signaling and phosphorylation of downstream kinases, including MAPK1/ERK2 and/or MAPK3/ERK1. Mutations that cause constitutive kinase activity promote cell proliferation and resistance to apoptosis via the activation of multiple signaling pathways.</p> |
| Usage | Research use only |
| Conjugate | PE-conjugated |
| DIMA Disclaimer | All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scr |

