Cat. No. DMC100274P



PRODUCT INFORMATION

Clone ID **DMC274**

IL5 **Target**

Synonyms IL-5;TRF;Interleukin-5;EDF

Host Species

PE-conjugated Anti-IL5 antibody(DMC274); IgG1 Description

Chimeric mAb

Delivery Under Development

Uniprot ID P05113

Rabbit/Human Fc chimeric IgG1 IgG type

Clonality Monoclonal Reactivity Human **Applications** Flow Cyt

Recommended

Flow Cyt 1:100 **Dilutions**

Purified from cell culture supernatant by affinity **Purification**

chromatography

Formulation & Reconstitution

Background

Usage

Liquid

□PBS with 0.05% Proclin300, 1% BSA

Storage & Shipping Store at 2°C-8°C for 6 months

> This gene encodes a cytokine that acts as a growth and differentiation factor for both B cells and eosinophils. The encoded cytokine plays a major role in the regulation of eosinophil formation; maturation; recruitment and survival. The increased production of this cytokine may be related to pathogenesis of eosinophil-dependent inflammatory diseases. This cytokine functions by binding to its receptor; which is a heterodimer; whose beta subunit is shared with the receptors

> for interleukine 3 (IL3) and colony stimulating factor 2 (CSF2:GM-CSF). This gene is located on chromosome 5 within a cytokine gene cluster which includes interleukin 4 (IL4); interleukin 13 (IL13); and CSF2 . This gene; IL4; and IL13 may be regulated coordinately by long-range regulatory elements spréad over 120 kilobases on

chromosome 5q31. Research use only

Conjugate PE-conjugated

All DIMA recombinant antibodies are genuinely

Email: info@dimabio.com Website: www.dimabio.com

generated by DIMA Biotech. They are all under patent application. Any protein sequencing or **DIMA Disclaimer** reverse engineering attempt is prohibited. We are

actively scr

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China)

/+86-400-006-0995(China)

