

PRODUCT INFORMATION

| | |
|---|--|
| Target | S1 protein NTD |
| Synonyms | S1 protein NTD; Spike protein S1 NTD; BetaCoV S1-NTD; COVID-19 |
| Description | Recombinant SARS-CoV-2 (2019-nCoV) S1 protein NTD with C-terminal 6×His tag |
| Delivery | In Stock |
| Uniprot ID | P0DTC2 |
| Expression Host | HEK293 |
| Tag | C-6×His Tag |
| Molecular Characterization | S1 protein NTD(Ser13-Leu303) 6×His tag |
| Molecular Weight | The protein has a predicted molecular mass of 33.7 kDa after removal of the signal peptide. The apparent molecular mass of S1-NTD-His is approximately 55 kDa due to glycosylation. |
| Purity | The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining. |
| Formulation & Reconstitution | Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. |
| Storage & Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | SARS-CoV-2 (Severe Acute Respiratory Syndrome Coronavirus 2) also known as Covid19 (2019 Novel Coronavirus) is a virus that causes illnesses ranging from the common cold to severe diseases. The spike protein is a type I transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor binding domain (RBD), which accounts for recognizing the cell surface receptor, ACE2. S2 contains basic elements needed for the membrane fusion. Recent publications indicate that S1-RBD domain can induce virus neutralizing-antibody and T cell response. |
| Usage | Research use only |
| Conjugate | Unconjugated |






Figure 1. SARS-CoV-2 (2019-nCoV) S1 protein NTD, His Tag on SDS-PAGE under reducing condition.

